

How to use Albendazole/Zentel/Alben and important safety information

For more information, please consult your doctor.

GSK encourages adverse events notifications in relation to its products at the following email pm.Safety@gsk.com

NX-BJ-ALN-PPT-260001/01-06-2026

How to use Albendazole/Zentel/ Alben tablets

Adults and children over two years of age:

For the treatment of intestinal parasites (worms)
take 400 mg as a single dose

= **one** Albendazole/Zentel/Alben **400 mg** tablet,
or

= **two** Albendazole/Zentel/Alben **200 mg** tablets

For all other situations – please consult your doctor.

How to use Albendazole/Zentel/Alben suspension

Adults and children over two years of age:

For the treatment of intestinal parasites (worms) take 400 mg as a single dose

= **20 ml** of the Albendazole/Zentel/Alben **2%** (200mg/10ml) suspension, or

= **10 ml** of the Albendazole/Zentel/Alben **4%** (400mg/10ml) suspension

For all other situations – please consult your doctor.

How to use Albendazole/Zentel/Alben suspension

Children from one to two years of age:

For the treatment of intestinal parasites (worms) take 200 mg as a single dose

= **10 ml** of the Albendazole/Zentel/Alben **2%** (200mg/10ml) suspension, or

= **5 ml** of the Albendazole/Zentel/Alben **4%** (400mg/10ml) suspension

For all other situations – please consult your doctor.

Important safety information (1)

Do not use Albendazole/Zentel/Alben if you are pregnant.

Women of childbearing age should only use Albendazole/Zentel/Alben during the first 10 days of starting their period, or immediately after a negative pregnancy test.

Important safety information (2)

Some people, particularly young children, may experience **problems swallowing Albendazole/Zentel/Alben tablets whole.**

To avoid any problems, the **tablets can be broken, crushed or chewed with a little water.** Alternatively, Albendazole/Zentel/Alben suspension can be used.

Important safety information (3)

If you have been **diagnosed with liver problems**, please consult a doctor before taking Albendazole/Zentel/Alben.

Please also consult a doctor if you experience **yellowing of the skin or eyes** after taking Albendazole/Zentel/Alben.

Important safety information (4)

Please consult a doctor as soon as possible if you develop **severe headaches with nausea and vomiting, fits and problems with your vision** after taking Albendazole/Zentel/Alben.

Be healthy!

Regarding the occurrence of adverse reactions after administration of the drug, please report to GlaxoSmithKline Export Limited by e-mail: pm.safety@gsk.com

Detailed information about this medicine is provided in the Patient Information Leaflet.

GlaxoSmithKline Export Limited, at 79 New Oxford Street, London, WC1A 1DG, UK.

©2026 GSK group of companies or its licensor.

Trademarks are owned by or licensed to the GSK group of companies.

